

Pathomechanisms and management of osteoporotic pain with no traumatic evidence

Sumihisa Orita, Kazuhide Inage, Miyako Suzuki, Kazuki Fujimoto, Kazuyo Yamauchi, Junichi Nakamura, Yusuke Matsuura, Takeo Furuya, Masao Koda, Kazuhisa Takahashi and Seiji Ohtori

Department of Orthopaedic Surgery, Graduate School of Medicine, Chiba University, Chiba, Japan

Abstract:

Introduction: Osteoporosis is a pathological state with an unbalanced bone metabolism mainly caused by accelerated osteoporotic osteoclast activity due to a postmenopausal estrogen deficiency, and it causes some kinds of pain, which can be divided into two types: traumatic pain due to a fragility fracture from impaired rigidity, and pain derived from an osteoporotic pathology without evidence of fracture. We aimed to review the concepts of osteoporosis-related pain and its management.

Methods: We reviewed clinical and basic articles on osteoporosis-related pain, especially with a focus on the mechanism of pain derived from an osteoporotic pathology (i.e., osteoporotic pain) and its pharmacological treatment.

Results: Osteoporosis-related pain tends to be robust and acute if it is due to fracture or collapse, whereas pathology-related osteoporotic pain is vague and dull. Non-traumatic osteoporotic pain can originate from an undetectable microfracture or structural change such as muscle fatigue in kyphotic patients. Furthermore, basic studies have shown that the osteoporotic state itself is related to pain or hyperalgesia with increased pain-related neuropeptide expression or acid-sensing channels in the local tissue and nervous system. Traditional treatment for osteoporotic pain potentially prevents possible fracture-induced pain by increasing bone mineral density and affecting related mediators such as osteoclasts and osteoblasts. The most common agent for osteoporotic pain management is a bisphosphonate. Other non-osteoporotic analgesic agents such as celecoxib have also been reported to have a suppressive effect on osteoporotic pain.

Conclusions: Osteoporotic pain has traumatic and non-traumatic factors. Anti-osteoporotic treatments are effective for osteoporotic pain, as they improve bone structure and the condition of the pain-related sensory nervous system. Physicians should always consider these matters when choosing a treatment strategy that would best benefit patients with osteoporotic pain.

Keywords:

osteoporotic pain, osteoporosis, anti-osteoporotic treatment

Spine Surg Relat Res 2017; 1(3): 121-128
dx.doi.org/10.22603/ssrr.1.2016-0001

Introduction

Chronic pain derived from musculoskeletal disorders plays a key role in the health profile of the general elderly population¹⁾, and it leads to impairments in quality of life (QOL) and in activities of daily living (ADLs), resulting in dependency, institutionalization, and increased health care costs²⁾. In particular, lower back pain (LBP) is one of the most important public health issues, as it is one of the most common symptoms that middle-aged and older people experience in their lives³⁾.

Among postmenopausal women, a significant increase in patients with LBP is often observed⁴⁾, and it is mainly due to osteoporosis, which is a pathological state with unbalanced bone metabolism chiefly caused by accelerated osteoporotic osteoclast activity due to a postmenopausal estrogen deficiency. Osteoporosis can cause pain from two main pathologies. First, pain can be due to a bone fracture, especially in the vertebrae and femoral neck because of their decreased bone rigidity (e.g., a fragility fracture). Its incidence according to the skeletal site is reported to be 27% of vertebral fracture, 19% of wrist, 14% of hip, 7% of pelvic, and

Corresponding author: Sumihisa Orita, sorita@chiba-u.jp

Received: September 30, 2016, Accepted: January 6, 2017

Copyright © 2017 The Japanese Society for Spine Surgery and Related Research

33% of other sites⁵). Vertebral fragility fractures are especially associated with significant rates of morbidity and mortality. This results in excess medical expenditures of \geq \$1 billion in the US⁶. Second, pain can originate from the osteoporotic state itself without any evidence of a fracture. A recent population-based survey of 3,097 older subjects (\geq 65 years old) reported that older adults with musculoskeletal complaints, including osteoporosis and chronic back pain, have considerable ADL problems⁷ due to pain.

Regarding the spinal osteoporosis itself, several kinds of osteoporotic agents are available: antiresorptive agents such as antiresorptives/denosumab, bone anabolic agents such as teriparatide, and other agents such as calcitonin, a selective estrogen receptor modulator (SERM), and Vitamin D. A previous systematic review shows the high evidence of preventing vertebral fractures in bisphosphonates, denosumab, and teriparatide with a relative risk reduction from 0.40 to 0.60⁸. However, the pathomechanisms and treatment for osteoporosis-related pain have not been adequately addressed yet.

The current article reviews osteoporosis-related pain, especially with a focus on LBP due to non-traumatic pain.

1) Traumatic osteoporosis-related pain with fracture or collapse

Osteoporotic patients tend to have fractures due to impaired bone mineral density (BMD). The main types of osteoporotic fractures are a (1) clinical fracture with a vertebral fracture or collapse with a distinguishing pain, including non-bony union or pseudoarthrosis, and a (2) morphometric fracture with gradual progression and little or almost no pain. Clinically, an osteoporosis-related fragility fracture is the main cause of LBP in elderly, with a 6%-8% incidence in women in their 60s and a \geq 30% incidence in those in their late 70s, and half of the individuals tend to have multiple vertebral fractures⁹. Some studies have reported that the 5-year survival rate is significantly lower in those with a fragility fracture of the vertebrae and femoral neck than in those with other fractures, which suggests the importance of treating these fractures to improve patients' ADL and QOL as well as control their pain¹⁰. Some of these fragile vertebral fractures result in a non-bony union (pseudoarthrosis) with more than a 9-month history⁵, which causes chronic back pain in the elderly. Though the actual incidence of the vertebral pseudoarthrosis is not clear, its incidence is estimated to amount to approximately 10% of the whole incidence of vertebral fragility fractures¹¹ followed by possible impairment in ADL and QOL. Toyone et al. has reported that the changes in vertebral wedging between standing and supine position are significantly associated with back pain¹². For those with pseudoarthrosis, analgesic agents such as NSAIDs and acetaminophen are administered^{13,14}. Some osteoporotic agents are reported to alleviate pain as is described later. Orthotic braces in the acute phase are reported to be effective in some systematic reviews, while its evidence level is relatively low¹³⁻¹⁶. When the pain is refrac-

tory to these kinds of conservative treatments, surgical treatments are available such as anterior decompression and stabilization¹⁷, osteotomy/shortening¹⁸, and percutaneous methods such as vertebroplasty and kyphoplasty¹⁹⁻²³. These surgeries are also important for preventing delayed neurologic deficit after vertebral arthrosis^{11,24,25}. However, a morphometric fracture is commonly observed in osteoporotic patients in their 60s and 70s with a progressive decrease in their height by about 1 cm yearly due to gradual progressive vertebral collapse²⁶. A previous study showed that pain from this kind of slowly progressing acute fracture can be diagnosed using magnetic resonance imaging, single photon emission computed tomography radioisotope, and bone scintigraphy in addition to radiographs²⁷.

2) Osteoporotic pain derived from an osteoporotic pathology

In contrast to patients with pain from a fragility fracture, some osteoporotic patients complain of vague LBP with discomfort without any evidence of a clinical fracture²⁸. The pain can be derived from multiple factors, including undetectable traumatic factors such as microfractures. Furthermore, in osteoporotic patients without any evidence of fractures, pain originates from an altered construction of bone and/or osteoporotic pathology. In patients with a degenerative hunched back, chronic LBP can be derived from deformed elements such as facet joints, cartilage, and ligament tissues surrounding the intervertebral discs as well as from muscle tissues and fascia that support the spine. Particularly, when the deformity of the lumbar spine is kyphotic, the erector spinae muscles become stretched, causing constant muscle contraction to cope with the condition, resulting in chronic muscle fatigue. These conditions induce increased intramuscular pressure with tense fascia, which result in an ischemic status in the muscle with chronic dull pain. Thus, these changes in the muscle cause LBP even for a short period (e.g., while standing and walking)²⁹⁻³⁶. The relationship between spinal kyphosis and the structural and functional parameters of trunk muscles should be investigated³⁶. A previous basic study suggested that the involvement of inflammation of the fascia in the back could alter the gait pattern in rodents³⁷.

In addition, the osteoporotic state itself is related to pain^{38,39}. Some basic studies have shown that the pathology itself induces specific hypersensitivity to pain, which is an unusual symptom of menopause^{40,41}. This kind of osteoporotic pain tends to have a chronic course, whereas traumatic osteoporotic pain tends to develop into acute pain. Herein, pain derived from osteoporosis without any fractures or injuries is defined as osteoporotic pain (Fig. 1).

The Mechanism of Osteoporotic Pain

Basic studies have shown evidence of osteoporotic pain using an osteoporotic animal model of ovariectomized (OVX) rats, which exhibit similar hormonal changes to that

of osteoporotic patients. Regarding pain, these osteoporotic models reportedly show hyperalgesia with significantly shortened latencies of tail withdrawal from hot water⁴²⁾ and long-term, formalin-induced licking⁴³⁾. The altered pain threshold in OVX rats can be caused by two anatomical factors: changes in the local tissue and changes in the nervous system.

The vertebral body itself is anatomically innervated by pain-related intravertebral sensory nerve fibers (Fig. 2)^{39,44)}, and nociceptors respond to mechanical, thermal, and chemical stimuli. Injury or inflammation results in the release of various chemical mediators (e.g., prostaglandins, cytokines, and growth factors). These chemical mediators stimulate osteoclast activity, activate nociceptors, and decrease their threshold for activation^{45,46)}. Additionally, transient-receptor potential vanilloid 1 (TRPV1) has also been reported to be upregulated in the dorsal root ganglion (DRG) of OVX rats³⁹⁾. TRPV1 is a ligand-gated, non-selective cation channel

preferentially expressed in pain-related, small-diameter primary afferent neurons^{47,48)}, and it responds to capsaicin and noxious heat and acid. Osteoporotic osteoclasts degrade bone minerals by secreting protons through the vacuolar H⁺-ATPase, creating local acidic microenvironments by inflammation^{49,50)}, which should evoke the stimulation of TRPV1. Furthermore, this acidic microenvironment produced by inflammation stimulates acid-sensing ion channels (ASICs) in acid-sensing nociceptors^{51,52)}. These stimulations evoked by osteoporotic osteoclasts lead to the chronic increased expression of inflammatory pain-related neuropeptides, calcitonin gene-related peptides (CGRP), and DRG neurons, innervating osteoporotic vertebrae³⁹⁾.

These changes in DRG can change the pain threshold at the corresponding levels of sensory distribution at the skin (dermatome) and bone (sclerotome). Fig. 3 demonstrates the distribution of the sclerotome of the lumbosacral and pelvic parts of rats, which explains the possible pain induced by pathological DRG⁵³⁾.

The activation of pain-related molecules upregulates the pain-related pathway at the more central level of the nervous system. In the dorsal horn of the spinal cord, c-Fos protein increases in response to pain stimuli from the peripheral primary afferent nervous system. c-Fos protein is a product of the c-Fos proto-oncogene expression, and it is expressed by noxious and non-noxious stimuli in the postsynaptic neurons of the dorsal horn of the spinal cord⁵⁴⁾. The reported increase in c-Fos in the spinal cord of OVX rats indicates the involvement of osteoporotic pain⁵⁵⁾. Additionally, an estrogen deficiency in osteoporotic patients causes bone loss and alters the spinal serotonergic system by suppressing serotonin (5-HT) receptor expression, which usually plays an important role at the dorsal horn of the spinal cord, resulting in hyperalgesia⁵⁶⁾. These findings such as increased production




Figure 1. A Venn diagram illustrating osteoporotic pain. Osteoporotic pain is derived from an osteoporotic pathology without fracture or injury.




Figure 2. Sensory innervation of a vertebral body (a) The vertebral body is innervated by sensory neurons that are mainly distributed from the posterior wall of vertebrae (adopted and modified from reference 46). (b) An immunofluorescent image of a vertebral section of an ovariectomized rat (merged and immunostained using Tuj-1 [a marker for nerve fibers], and a calcitonin gene-related peptide [an inflammatory pain-related neuropeptide]) (adopted and modified from reference 39). Arrowheads indicate that the sensory fiber innervating the bone marrow (BM). TB: trabecular bone




Figure 3. Scheme showing the sclerotome of rats. A: Anteroposterior view, B: lateral view. Segmental distributions of sensory innervation in bones indicate possible reflexion of the altered property of the corresponding sensory neurons (adopted and modified from reference 53).

of pain-related channels, receptors, and neuropeptides in DRG and spinal cord as well as change in the local tissue are included in the mechanism of osteoporotic pain.

In addition to these static changes in osteoporotic vertebrae, other dynamic factors can affect osteoporotic pain, e. g., gravity, especially the axial load on the vertebral axis. A previous basic study simulated the axial load using coccygeal vertebrae that were continuously compressed with experimental rubber bands⁵⁷⁾ (Fig. 4) to demonstrate the upregulated expression of a neurological injury-related neuropeptide, activating transcription factor 3, as well as CGRPs in the DRG innervating the vertebrae. The authors reported an increased expression of both peptides, so they concluded that osteoporotic pain is caused by a neuropathic factor as well as an inflammatory (nociceptive) one. The result of the study indicates the complicated and chronic mechanism of osteoporotic pain.

Treatment for Osteoporotic Pain

Traditional osteoporosis treatment for pain potentially includes the prevention of possible fracture-induced pain by increasing BMD, which each agent initially aims to achieve. Moreover, each anti-osteoporosis agent has been reported to have a specific pain-related active site, which will be described further in the following sections (Fig. 5).




Figure 4. An experiment using the coccygeal vertebrae to investigate the effect of the axial load on sensory innervation (adopted and modified from reference 57). Co: coccygeal, FG: Fluorogold (a retrograde neurotracer) applied onto the dorsal periosteal surface of the Co5 vertebra. The 0.8-mm (diameter) k-wires penetrated the Co4 and 6 vertebrae, and they were connected using experimental rubber bands with a continuous compressive load.

Antiresorptive agents: Bisphosphonate and denosumab

A bisphosphonate is the most common anti-osteoporosis agent that inhibits osteoclast activity to improve BMD^{58,59)}. In addition, there have been some reports of using bisphosphonates for pain.

For patients with an osteoporotic fragility fracture, intravenous pamidronate is effective for chronic back pain⁶⁰⁾. Ibandronate is reported to suppress the expression of substance P mRNA and tumor necrosis factor- α in the DRG of a rat model with persistent inflammation⁶¹⁾. Another bisphosphonate, risedronate, has been reported to suppress the expression of CGRP in the DRG³⁸⁾ to improve LBP in clinical postmenopausal osteoporotic patients without any evidences of vertebral fractures³⁸⁾. Furthermore, an *in vitro* experiment showed the inhibitory effect of risedronate on axonal growth of neurite of pain-related DRG neurons from rat neonates³⁹⁾, indicating that a bisphosphonate itself can produce an analgesic effect by suppressing peripheral nerve function. Moreover, a third-generation of minodronic acid has been reported to have an analgesic effect by antagonizing purinergic P2X(2/3) receptor function⁶²⁾. One study showed that pa-




Figure 5. Scheme of osteoporotic pain and the active sites of anti-osteoporotic agents: 5-HT: hydroxytryptamine (serotonin), ASIC: acid-sensing ion channel, TRPV1: transient-receptor potential vanilloid 1, TNF- α : tumor necrosis factor- α , SERM: selective estrogen receptor modulator, PTH: parathyroid hormone

tients treated with daily minodronate showed more pain relief measured on the visual analog scale⁶³.

A bisphosphonate has effects on other molecules too. Alendronate improves osteoporotic pain by inhibiting TRPV1⁵². In addition, a randomized, controlled trial showed that alendronate produced a stronger analgesic effect than calcitonin in postmenopausal osteoporotic women⁶⁴. Furthermore, combination therapy using risedronate and treadmill exercise in OVX rats showed effective improvement in BMD and decreased expression of pain-related neuropeptides³⁹, which proves the effect of physical exercise as anti-osteoporotic treatment⁶⁵⁻⁶⁸.

Recently, another type of anti-osteoporotic agent, denosumab (an anti-receptor activator of nuclear factor- κ B ligand [RANKL] antibody), has been discussed. It has a strong suppressive effect on osteoclasts by inhibiting their differentiation from precursors, and it has a significant effect on BMD improvement^{69,70}. RANKL itself is involved in pain pathology⁷¹, and an antagonism against RANKL with denosumab has been reported to improve LBP in patients with osteoporotic pain⁷².

Estrogen and Selective Estrogen Receptor Modulator

Administering estrogen or a selective estrogen receptor

modulator (SERM) seems reasonable for treating osteoporosis, because the pathology is based on decreased estrogen secretion, which suppresses CGRP expression of DRG⁷³.

However, SERM has estrogen and estrogen-like effects on bone, and it antagonizes the action of estrogen in the endometrium and breast tissue. A previous study reported that treatment with raloxifene in postmenopausal osteoporotic women showed a marked reduction in musculoskeletal pain⁷⁴.

Calcitonin

Calcitonin is an endogenous polypeptide hormone that inhibits bone resorption by osteoclasts⁷⁵. It decreases hyperalgesia in OVX rats by upregulating the activity of the descending serotonergic inhibitory system⁷⁶ by recovering the resumption of the synthesis of 5-HT receptors^{5,6,77}. Another clinical study reported that it has an analgesic effect comparative to morphine⁷⁸ with significantly increased plasma β -endorphin levels in patients with postmenopausal osteoporosis⁷⁹. Furthermore, a clinical study indicated that the use of elcatonin better reduced chronic back pain when combined with risedronate for more than 3 months⁸⁰. These facts prove the analgesic effect of calcitonin on osteoporotic pain.

Vitamin D

Vitamin D is reported to stimulate osteoblast activity and other effects such as an increase in the intestinal absorption of calcium, and regulation of parathyroid hormone activity, calcium metabolism, and proximal muscle function⁸¹. Vitamin D-related pain pathways are associated with changes in cortical, hormonal, neuronal, and immunological alterations⁸². Vitamin D alleviates musculoskeletal pain and down-regulates inflammatory markers in patients with osteoporosis, especially in those treated with zoledronic acid⁸³.

Teriparatide

Teriparatide is a human parathyroid hormone analog that induces the formation of new bone by increasing osteoblast generation and stimulating osteoclasts to resorb bone. It significantly induces increased BMD and lowers the incidence of new fractures to achieve improved QOL⁸⁴. Its analgesic effect on osteoporosis-related pain includes a strengthened bone structure, which results in a lower incidence of new microfracture and macrofractures⁸⁵. In a clinical study, teriparatide is reported to significantly reduce low back pain better than risedronate⁸⁶. Because it is a relatively new agent, there is still a lack of evidence using teriparatide to relieve osteoporotic pain, thus this should be investigated in a future study.

Other non-osteoporotic agents

Other non-osteoporotic analgesic agents reportedly have a suppressive effect on osteoporotic pain. Non-steroidal anti-inflammatory drugs (NSAIDs) and opioids are widely used for chronic pain associated with fragility or traumatic fractures. Additionally, some NSAIDs such as celecoxib and cyclooxygenase-2 selective agents suppress osteoporotic pain by inhibiting carbonic anhydrase II in the osteoclasts, which indicates a possible analgesic effect on osteoporotic pain.

Considering the aforementioned information, anti-osteoporotic treatments are effective for relieving osteoporotic pain by improving bone structure and the condition of the pain-related sensory nervous system.

Conclusions

Osteoporotic pain derived from an osteoporotic pathology is a condition induced by an increase in pain-related neuropeptides in the sensory neurons innervating osteoporotic vertebrae, where activated osteoclasts have more inflammatory conditions that stimulate pain-inducing molecules with acidic inflammatory circumstances. Osteoporosis treatment predominantly aims to increase the BMD of patients to prevent possible fragility fractures, and recent considerable evidence shows that anti-osteoporotic treatment also relieves pain. Physicians should always consider these matters when choosing a treatment strategy that would best benefit patients with osteoporotic pain.

Conflicts of Interest: The authors declare that there are no conflicts of interest.

Sources of funding: none

References

1. Badley EM, Rasooly I, Webster GK. Relative importance of musculoskeletal disorders as a cause of chronic health problems, disability, and health care utilization: findings from the 1990 Ontario Health Survey. *J Rheumatol*. 1994;21(3):505-14.
2. Woolf AD, Erwin J, March L. The need to address the burden of musculoskeletal conditions. *Best Pract Res Clin Rheumatol*. 2012; 26(2):183-224.
3. Cassidy JD, Carroll LJ, Cote P. The Saskatchewan health and back pain survey. The prevalence of low back pain and related disability in Saskatchewan adults. *Spine*. 1998;23(17):1860-6;discussion 7.
4. Adera T, Deyo RA, Donatelle RJ. Premature menopause and low back pain. A population-based study. *Ann Epidemiol*. 1994;4(5): 416-22.
5. Burge R, Dawson-Hughes B, Solomon DH, et al. Incidence and economic burden of osteoporosis-related fractures in the United States, 2005-2025. *J Bone Miner Res*. 2007;22(3):465-75.eng.
6. Goldstein CL, Chutkan NB, Choma TJ, et al. Management of the elderly with vertebral compression fractures. *Neurosurgery*. 2015; 77 Suppl 4:S33-45.
7. Stamm TA, Pieber K, Crevenna R, et al. Impairment in the activities of daily living in older adults with and without osteoporosis, osteoarthritis and chronic back pain: A secondary analysis of population-based health survey data. *BMC Musculoskelet Disord*. 2016;17:139.eng.
8. Crandall CJ, Newberry SJ, Diamant A, et al. Comparative effectiveness of pharmacologic treatments to prevent fractures: an updated systematic review. *Ann Intern Med*. 2014;161(10):711-23.
9. Ross PD, Fujiwara S, Huang C, et al. Vertebral fracture prevalence in women in Hiroshima compared to Caucasians or Japanese in the US. *Int J Epidemiol* 1995;24(6):1171-7.
10. Johnell O, Kanis JA, Oden A, et al. Mortality after osteoporotic fractures. *Osteoporosis Int*. 2004;15(1):38-42.
11. Pappou IP, Papadopoulos EC, Swanson AN, et al. Osteoporotic vertebral fractures and collapse with intravertebral vacuum sign (Kummel's disease). *Orthopedics*. 2008;31(1):61-6.eng.
12. Toyone T, Toyone T, Tanaka T, et al. Changes in vertebral wedging rate between supine and standing position and its association with back pain: a prospective study in patients with osteoporotic vertebral compression fractures. *Spine* 2006;31(25):2963-6.
13. Vorsanger GJ, Farrell J, Xiang J, et al. Tapentadol, oxycodone or placebo for acute pain of vertebral compression fractures: a randomized Phase IIIb study. *Pain Manag*. 2013;3(2):109-18.
14. Rzewuska M, Ferreira M, McLachlan AJ, et al. The efficacy of conservative treatment of osteoporotic compression fractures on acute pain relief: a systematic review with meta-analysis. *Eur Spine J*. 2015;24(4):702-14.
15. Pfeifer M, Kohlwey L, Begerow B, et al. Effects of two newly developed spinal orthoses on trunk muscle strength, posture, and quality-of-life in women with postmenopausal osteoporosis: a randomized trial. *Am J Phys Med Rehabil*. 2011;90(10):805-15.
16. Pfeifer M, Begerow B, Minne HW. Effects of a new spinal orthosis on posture, trunk strength, and quality of life in women with postmenopausal osteoporosis: a randomized trial. *Am J Phys Med Rehabil*. 2004;83(3):177-86.
17. Kaneda K, Taneichi H, Abumi K, et al. Anterior decompression and stabilization with the Kaneda device for thoracolumbar burst fractures associated with neurological deficits. *J Bone Joint Surg*

- Am. 1997;79(1):69-83.
18. Saita K, Hoshino Y, Kikkawa I, et al. Posterior spinal shortening for paraplegia after vertebral collapse caused by osteoporosis. *Spine*. 2000;25(21):2832-5.
 19. Kim DY, Lee SH, Jang JS, et al. Intravertebral vacuum phenomenon in osteoporotic compression fracture: report of 67 cases with quantitative evaluation of intravertebral instability. *J Neurosurg*. 2004;100(1 Suppl Spine):24-31.eng.
 20. Ishiguro S, Kasai Y, Sudo A, et al. Percutaneous vertebroplasty for osteoporotic compression fractures using calcium phosphate cement. *J Orthop Surg. (Hong Kong)*. 2010;18(3):346-51.eng.
 21. Nakano M, Hirano N, Zukawa M, et al. Vertebroplasty Using Calcium Phosphate Cement For Osteoporotic Vertebral Fractures: study of outcomes at a minimum follow-up of two years. *Asian Spine J*. 2012;6(1):34-42.eng.
 22. Nakano M, Hirano N, Matsuura K, et al. Percutaneous transpedicular vertebroplasty with calcium phosphate cement in the treatment of osteoporotic vertebral compression and burst fractures. *J Neurosurg*. 2002;97(3 Suppl):287-93.eng.
 23. Yang H, Wang G, Liu J, et al. Balloon kyphoplasty in the treatment of osteoporotic vertebral compression fracture nonunion. *Orthopedics*. 2010;33(1):24.eng.
 24. Uchida K, Kobayashi S, Matsuzaki M, et al. Anterior versus posterior surgery for osteoporotic vertebral collapse with neurological deficit in the thoracolumbar spine. *Eur Spine J*. 2006;15(12):1759-67.eng.
 25. Sasaki Y, Aoki Y, Nakajima A, et al. Delayed neurologic deficit due to foraminal stenosis following osteoporotic late collapse of a lumbar spine vertebral body. *Case Rep Orthop*. 2013;2013:682075.eng.
 26. Neer RM, Arnaud CD, Zanchetta JR, et al. Effect of parathyroid hormone (1-34) on fractures and bone mineral density in postmenopausal women with osteoporosis. *N Engl J Med*. 2001;344(19):1434-41.
 27. Ryan PJ, Evans P, Gibson T, et al. Osteoporosis and chronic back pain: A study with single-photon emission computed tomography bone scintigraphy. *J Bone Miner Res*. 1992;7(12):1455-60.
 28. Nakamura T. Low back pain accompanying osteoporosis. *Japan Med Assoc J*. 2003;46(10):445-51.
 29. Itoi E. Roentgenographic analysis of posture in spinal osteoporotics. *Spine* 1991;16(7):750-6.
 30. Miyakoshi N, Itoi E, Kobayashi M, et al. Impact of postural deformities and spinal mobility on quality of life in postmenopausal osteoporosis. *Osteoporos Int*. 2003;14(12):1007-12.
 31. Ranger TA, Teichtahl AJ, Cicuttini FM, et al. Shorter Lumbar Paraspinal Fascia Is Associated With High Intensity Low Back Pain and Disability. *Spine*. 2016;41(8):E489-93.eng.
 32. Yabuki S, Kikuchi S, Midorikawa H, et al. Vascular backache and consideration of its pathomechanisms: report of two cases. *J Spinal Disord*. 1999;12(2):162-7.eng.
 33. Pfeifer M, Sinaki M, Geusens P, et al. Musculoskeletal rehabilitation in osteoporosis: a review. *J Bone Miner Res*. 2004;19(8):1208-14.eng.
 34. Handa N, Yamamoto H, Tani T, et al. The effect of trunk muscle exercises in patients over 40 years of age with chronic low back pain. *J Orthop Sci*. 2000;5(3):210-6.eng.
 35. Takemasa R, Yamamoto H, Tani T. Trunk muscle strength in and effect of trunk muscle exercises for patients with chronic low back pain. The differences in patients with and without organic lumbar lesions. *Spine*. 1995;20(23):2522-30.eng.
 36. Roghani T, Zavieh MK, Manshadi FD, et al. Age-related hyperkyphosis: update of its potential causes and clinical impacts-narrative review. *Aging Clin Exp Res*. 2016.eng.
 37. Miyagi M, Ishikawa T, Kamoda H, et al. Assessment of gait in a rat model of myofascial inflammation using the CatWalk system. *Spine (Phila Pa 1976)*. 2011;36(21):1760-4.
 38. Ohtori S, Akazawa T, Murata Y, et al. Risedronate decreases bone resorption and improves low back pain in postmenopausal osteoporosis patients without vertebral fractures. *J Clin Neurosci*. 2010;17(2):209-13.eng.
 39. Orita S, Ohtori S, Koshi T, et al. The effects of risedronate and exercise on osteoporotic lumbar rat vertebrae and their sensory innervation. *Spine*. 2010;35(22):1974-82.
 40. Yeh CB, Weng SJ, Chang KW, et al. Calcitonin alleviates hyperalgesia in osteoporotic rats by modulating serotonin transporter activity. *Osteoporos Int*. 2016.eng.
 41. Sanoja R, Cervero F. Estrogen modulation of ovariectomy-induced hyperalgesia in adult mice. *Eur J Pain*. 2008;12(5):573-81.
 42. Forman LJ, Tingle V, Estilow S, et al. The response to analgesia testing is affected by gonadal steroids in the rat. *Life Sci*. 1989;45(5):447-54.
 43. Franceschini R, Bottaro P, Panopoulos C. Long-term treatment with salmon calcitonin in postmenopausal osteoporosis. *Curr Ther Res*. 1983;34:795-800.
 44. Bailey JF, Liebenberg E, Degmetich S, et al. Innervation patterns of PGP 9.5-positive nerve fibers within the human lumbar vertebra. *J Anat*. 2011;218(3):263-70.
 45. Haegerstam GA. Pathophysiology of bone pain: a review. *Acta orthopaedica Scandinavica*. 2001;72(3):308-17.
 46. Payne R. Mechanisms and management of bone pain. *Cancer*. 1997;80(8 Suppl):1608-13.
 47. Tominaga M, Caterina MJ. Thermosensation and pain. *J Neurobiol*. 2004;61(1):3-12.eng.
 48. Tominaga M, Caterina MJ, MalMBERG AB, et al. The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron*. 1998;21(3):531-43.
 49. Rousselle a-V, Heymann D. Osteoclastic acidification pathways during bone resorption. *Bone*. 2002;30(4):533-40.
 50. Teitelbaum SL. Bone Resorption by Osteoclasts. *Science*. 2000;289(5484):1504-8.
 51. Kanaya K, Iba K, Abe Y, et al. Acid-sensing ion channel 3 or P2X 2/3 is involved in the pain-like behavior under a high bone turnover state in ovariectomized mice. *J Orthop Res*. 2016;34(4):566-73.eng.
 52. Abe Y, Iba K, Sasaki K, et al. Inhibitory effect of bisphosphonate on osteoclast function contributes to improved skeletal pain in ovariectomized mice. *J Bone Miner Metab*. 2015;33(2):125-34.eng.
 53. Takahashi Y, Ohtori S, Takahashi K. Sclerotomes in the thoracic and lumbar spine, pelvis, and hindlimb bones of rats. *J Pain*. 2010;11(7):652-62.
 54. Hunt SP, Pini A, Evan G. Induction of c-fos-like protein in spinal cord neurons following sensory stimulation. *Nature*. 1987;328(6131):632-4.
 55. Lu Y, Li Z, Li HJ, et al. A comparative study of the effect of 17 beta-estradiol and estriol on peripheral pain behavior in rats. *Steroids*. 2012;77(3):241-9.
 56. Ito A, Kumamoto E, Takeda M, et al. Mechanisms for ovariectomy-induced hyperalgesia and its relief by calcitonin: participation of 5-HT1A-like receptor on C-afferent terminals in substantia gelatinosa of the rat spinal cord. *J Neurosci*. 2000;20(16):6302-8.
 57. Suzuki M, Orita S, Miyagi M, et al. Vertebral compression exacerbates osteoporotic pain in an ovariectomy-induced osteoporosis rat

- model. *Spine*. 2013;38(24):2085-91.eng.
58. Black DM, Thompson DE, Bauer DC, et al. Fracture risk reduction with alendronate in women with osteoporosis: the Fracture Intervention Trial. FIT Research Group. *J Clin Endocrinol Metab*. 2000;85(11):4118-24.
 59. Rogers MJ, Crockett JC, Coxon FP, et al. Biochemical and molecular mechanisms of action of bisphosphonates. *Bone*. 2011;49(1):34-41.
 60. Gangji V, Appelboom T. Analgesic effect of intravenous pamidronate on chronic back pain due to osteoporotic vertebral fractures. *Clin Rheumatol*. 1999;18(3):266-7.
 61. Bianchi M, Franchi S, Ferrario P, et al. Effects of the bisphosphonate ibandronate on hyperalgesia, substance P, and cytokine levels in a rat model of persistent inflammatory pain. *Eur J Pain*. 2008;12(3):284-92.
 62. Kakimoto S, Nagakura Y, Tamura S, et al. Minodronic acid, a third-generation bisphosphonate, antagonizes purinergic P2X(2/3) receptor function and exerts an analgesic effect in pain models. *Eur J Pharmacol*. 2008;589(1-3):98-101.
 63. Yoshioka T, Okimoto N, Okamoto K, et al. A comparative study of the effects of daily minodronate and weekly alendronate on upper gastrointestinal symptoms, bone resorption, and back pain in postmenopausal osteoporosis patients. *J Bone Miner Metab*. 2013;31(2):153-60.
 64. Iwamoto J, Makita K, Sato Y, et al. Alendronate is more effective than elcatonin in improving pain and quality of life in postmenopausal women with osteoporosis. *Osteoporos Int*. 2010.
 65. Iwamoto J, Sato Y, Takeda T, et al. Effectiveness of exercise in the treatment of lumbar spinal stenosis, knee osteoarthritis, and osteoporosis. *Aging Clin Exp Res*. 2010;22(2):116-22.eng.
 66. Iwamoto J, Takeda T, Sato Y. Effect of treadmill exercise on bone mass in female rats. *Exp Anim*. 2005;54(1):1-6.
 67. Li WC, Chen YC, Yang RS, et al. Effects of exercise programmes on quality of life in osteoporotic and osteopenic postmenopausal women: a systematic review and meta-analysis. *Clin Rehabil*. 2009;23(10):888-96.
 68. Sinaki M, Pfeifer M, Preisinger E, et al. The role of exercise in the treatment of osteoporosis. *Curr Osteoporos Rep*. 2010;8(3):138-44.eng.
 69. Bridgeman MB, Pathak R. Denosumab for the reduction of bone loss in postmenopausal osteoporosis: a review. *Clin Ther*. 2011;33(11):1547-59.eng.
 70. Petranova T, Sheytanov I, Monov S, et al. Denosumab improves bone mineral density and microarchitecture and reduces bone pain in women with osteoporosis with and without glucocorticoid treatment. *Biotechnol Biotechnol Equip*. 2014;28(6):1127-37.eng.
 71. Matsuyama Y, Sakuma Y, Suzuki M, et al. Evaluation of behavior and expression of receptor activator of nuclear factor-kappa B ligand in dorsal root Ganglia after sciatic nerve compression and application of nucleus pulposus in rats. *Asian Spine J*. 2014;8(5):557-64.
 72. Inage K, Orita S, Yamauchi K, et al. The time course changes in bone metabolic markers after administering the anti-receptor activator of nuclear factor-kappa b ligand antibody and drug compliance among patients with osteoporosis. *Asian Spine J*. 2015;9(3):338-43.
 73. Yang Y, Ozawa H, Lu H, et al. Immunocytochemical analysis of sex differences in calcitonin gene-related peptide in the rat dorsal root ganglion, with special reference to estrogen and its receptor. *Brain Res*. 1998;791(1-2):35-42.
 74. Scharla S, Oertel H, Helsing K, et al. Skeletal pain in postmenopausal women with osteoporosis: prevalence and course during raloxifene treatment in a prospective observational study of 6 months duration. *Curr Med Res Opin*. 2006;22(12):2393-402.
 75. Cranney A, Tugwell P, Zytaruk N, et al. Meta-analyses of therapies for postmenopausal osteoporosis. VI. Meta-analysis of calcitonin for the treatment of postmenopausal osteoporosis. *Endocr Rev*. 2002;23(4):540-51.
 76. Takayama B, Kikuchi Si, Konno Si, et al. An immunohistochemical study of the antinociceptive effect of calcitonin in ovariectomized rats. *BMC Musculoskelet Disord*. 2008;9:164-.
 77. Sun RQ, Tu YJ, Lawand NB, et al. Calcitonin gene-related peptide receptor activation produces PKA- and PKC-dependent mechanical hyperalgesia and central sensitization. *J Neurophysiol*. 2004;92(5):2859-66.
 78. Martin MI, Goicoechea C, Ormazabal MJ, et al. Analgesic effect of two calcitonins and in vitro interaction with opioids. *Gen Pharmacol*. 1995;26(3):641-7.
 79. Ofluoglu D, Akyuz G, Unay O, et al. The effect of calcitonin on beta-endorphin levels in postmenopausal osteoporotic patients with back pain. *Clin Rheumatol*. 2007;26(1):44-9.
 80. Hongo M, Miyakoshi N, Kasukawa Y, et al. Additive effect of elcatonin to risedronate for chronic back pain and quality of life in postmenopausal women with osteoporosis: a randomized controlled trial. *J Bone Miner Metab*. 2015;33(4):432-9.
 81. Paolucci T, Saraceni VM, Piccinini G. Management of chronic pain in osteoporosis: challenges and solutions. *J Pain Res* 2016;9:177-86.eng.
 82. Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc*. 2003;78(12):1463-70.
 83. Catalano A, Morabito N, Atteritano M, et al. Vitamin D reduces musculoskeletal pain after infusion of zoledronic acid for postmenopausal osteoporosis. *Calcif Tissue Int*. 2012;90(4):279-85.
 84. Manuele S, Sorbello L, Puglisi N, et al. The teriparatide in the treatment of severe senile osteoporosis. *Arch Gerontol Geriatr*. 2007;44 Suppl 1:249-58.
 85. Iolascon G, Gimigliano F, Malavolta N, et al. Effectiveness of teriparatide treatment on back pain-related functional limitations in individuals affected by severe osteoporosis: a prospective pilot study. *Clin Cases Miner Bone Metab*. 2012;9(3):161-5.
 86. Tsuchie H, Miyakoshi N, Kasukawa Y, et al. The effect of teriparatide to alleviate pain and to prevent vertebral collapse after fresh osteoporotic vertebral fracture. *J Bone Miner Metab*. 2016;34(1):86-91.

Spine Surgery and Related Research is an Open Access article distributed under the Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).